

Passive Muscle Stiffness of Biceps Femoris is Acutely Reduced after Eccentric Knee Flexion

メタデータ	言語: English 出版者: 公開日: 2023-07-07 キーワード (Ja): キーワード (En): 作成者: 支, 磊 メールアドレス: 所属:
URL	https://jair.repo.nii.ac.jp/records/2003322

Research article

Passive Muscle Stiffness of Biceps Femoris is Acutely Reduced after Eccentric Knee Flexion

Lei Zhi¹, Naokazu Miyamoto^{1,2}✉ and Hisashi Naito^{1,3}

¹ Graduate School of Health and Sports Science, Juntendo University, Inzai, Japan; ² Department of Sport Science and Research, Japan Institute of Sports Sciences, Tokyo, Japan; ³ Institute of Health and Sport Science & Medicine, Juntendo University, Inzai, Japan

Abstract

Eccentric hamstring exercises reportedly prevent hamstring strain injury in the biceps femoris long head (BFLh). However, information on the favorable adaptive responses in the BFLh to eccentric hamstring exercises is limited. We aimed to examine the acute effect of maximal isokinetic eccentric knee flexion on passive BFLh stiffness as a potential risk factor for the hamstring strain injury using ultrasound shear wave elastography. Ten young participants randomly performed both tasks involving five consecutive repetitions of isokinetic concentric and eccentric knee flexion with maximal effort on different legs. Passive BFLh shear modulus was taken before and 30, 60, 90, and 120 s after each task. Passive BFLh shear modulus was significantly reduced at all time points after eccentric knee flexion, whereas there was no significant change in passive BFLh shear modulus after the concentric task. The present findings indicate that passive BFLh stiffness would reduce specifically after low-volume, slow-velocity eccentric knee flexion exercise. The findings may help provide practitioners with a basis to develop more effective exercise programs for preventing HSI.

Key words: Eccentric hamstring exercises, isokinetic knee flexion, ultrasound shear wave elastography, shear modulus.

Introduction

Hamstring strain injury (HSI) remains one of the most common injuries across a range of sprint-based sports, such as track-and-field (Opar et al., 2014), soccer (Ekstrand et al., 2011), rugby (Brooks et al., 2005), and baseball (Ahmad et al., 2014). Eccentric hamstring exercises (i.e., lengthening of the contracting hamstring) reportedly prevent HSI (van Dyk et al., 2019) through possible mechanisms of increasing hamstring strength (Mjolsnes et al., 2004) and consequently hamstring-to-quadriceps strength ratio (Severo-Silveira et al., 2021) and/or of increasing lower limb flexibility (O'Sullivan et al., 2012). On the other hand, although most of the sprint-type HSI occurs in the long head of the biceps femoris (BFLh) (Askling et al., 2007), available information on the favorable adaptive responses in the BFLh to eccentric hamstring exercises is limited to an increase in BFLh fascicle length (Gerard et al., 2020) which has been suggested as a possible mechanism for the prevention of HSI (Timmins et al., 2016). A more detailed elucidation of the favorable adaptive responses of eccentric hamstring exercises on the BFLh (rather than the

hamstring) can lead to advanced strategies aimed at preventing HSI.

Stiffness of the hamstring muscle-tendon unit has been reported as a risk factor for HSI (Green et al., 2020; Watsford et al., 2010). Muscle-tendon unit stiffness is attributable, at least in part, to muscle stiffness (Gajdosik, 2001; Konrad and Tilp, 2020). Recently, high passive muscle stiffness measured with ultrasound shear wave elastography (SWE), which can quantify the stiffness of individual muscles, has been suggested as a factor related to muscle injury (Kumagai et al., 2019; Miyamoto-Mikami et al., 2021). Considering these observations, together with the fact that most HSI occurs in the BFLh as mentioned above (Askling et al., 2007), it is plausible to suppose that passive BFLh stiffness is a risk factor for HSI and that eccentric hamstring exercises prevent HSI by reducing passive BFLh stiffness. However, this is only speculation and not experimentally demonstrated. Therefore, it is important to pursue an understanding of whether and how eccentric hamstring exercise modifies passive BFLh stiffness.

As a first step toward the above goal, based on the notions that findings about the acute changes by exercises could provide insights into their chronic responses, the present study aimed to examine an acute effect of maximal isokinetic eccentric knee flexion on passive BFLh stiffness, with special emphasis on the comparison with concentric (i.e., shortening) contraction. Passive BFLh stiffness is reduced by hamstring stretching of passive knee extension (i.e., lengthening of the resting hamstring) in the hip flexed position (Miyamoto et al., 2017). Based on this observation, we hypothesized that passive BFLh stiffness would be acutely reduced by eccentric but not concentric knee flexion exercise.

Methods

Study design

Two tasks that consisted of isokinetic concentric (CON) and eccentric (ECC) knee flexion were performed to fulfill the goal of the present study. Except for the contraction type of hamstring exercise, the protocols and variables for analyses were the same between the two tasks (see below sections). In a randomized design, each participant performed a CON task on one leg and an ECC task on the other leg. The CON and ECC tasks were performed on the same day with a 10-min rest period in a randomized order. The

room temperature was set to 24°C using an air conditioner to minimize the potential temperature-induced effects.

Participants

When calculating the necessary sample size, there was no direct data from previous studies to be referred to for a priori power analysis since no previous study has investigated the acute effect of eccentric contractions on muscle stiffness. Consequently, the necessary sample size was calculated based on the data of our preliminary study [$n = 5$, partial η^2 for TASK (2) \times TIME (5) interaction = 0.689, partial η^2 for main effect of TIME in ECC task = 0.630], using a priori power analysis (G*Power, version 3.1.9.4 software, 26 Heinrich-Heine-Universität Düsseldorf, Germany) with an assumed type 1 error of 0.05 and statistical power ($1-\beta$) of 0.8. The critical sample size was estimated to be at least 10 for each task. Ten healthy men (age: 24.1 ± 1.8 years, height: 173.2 ± 5.0 cm, weight: 64.9 ± 4.8 kg; mean \pm SD) were recruited for the study. Participants with a history of thigh muscle or knee injuries within the last 6 months were excluded from the study. None of the participants had been involved in any resistance training exercise. They were asked to refrain from intense exercise within 48 h before the experiment. Before testing, the participants were informed of the study procedures and provided written informed consent. The study was approved by the local ethics committee (project identification code: 2021-96), and all experimental procedures were performed in accordance with the Declaration of Helsinki.

Experimental setup and procedure

Each participant was seated on an isokinetic dynamometer (Biodex System 4 Pro, Biodex Medical System, USA) with the hip flexed at 70° (0° = lying position). The participant was tightly secured to the dynamometer using non-elastic straps. The rotation axis of the motor was aligned with that of the knee joint. The lever arm of the dynamometer was attached 3–4 cm above the lateral malleolus. After a 5-min rest, passive BFlh shear modulus (a measure of stiffness, expressed in the unit of Pascal) was measured using ultrasound SWE (see below section) before isokinetic exercises (PRE) in the sitting position with the hip flexed at 70° and

the knee of the tested leg fully extended. Subsequently, the participants performed a specific isokinetic warm-up exercise consisting of 6 sub-maximal (3 self-perceived 50% effort and 3 self-perceived 80% effort) and 2 maximal CON or ECC knee flexion on the tested leg at an angular velocity of 20°/s to familiarize themselves with the protocol (El-Ashker et al., 2019). This angular velocity was chosen based on the fact that the Nordic hamstring exercise, one of the major exercises of HSI prevention programs, was performed at approximately 20°/s (Alt et al., 2018). After a 3-min rest, each participant performed five consecutive repetitions of CON or ECC knee flexion with maximal efforts. In both tasks, the knee joint range of motion was between 0° and 90° (0° = full extension). Measurements of passive BFlh shear modulus were taken at 30, 60, 90, and 120 s after the end of the CON/ECC task (POST).

SWE measurement

An ultrasound SWE scanner (Aixplorer Ver. 12; Supersonic Imagine, France), coupled with a linear array probe (SL10-2), was used to assess passive BFlh shear modulus in SWE mode (MSK preset, persistence = off, smoothing = 5). According to a previous study (Miyamoto et al., 2018), the ultrasound probe was positioned at 50% of the thigh length (the distance between the greater trochanter and the lateral epicondyle of the femur). The probe orientation was adjusted to identify fascicles within the B-mode image, and the location was marked on the skin with a waterproof pen. At each of five measurement time points (i.e., PRE, POST30, POST60, POST90, POST120) in each task, SWE measurements were performed three times (i.e., three images were acquired). The images were acquired while ensuring that the color map was stable for a few seconds. Care was taken not to press and deform the muscle while scanning. A region of interest in the SWE color map was carefully selected as large as possible with the exclusion of subcutaneous tissues and aponeurosis for each image using the built-in software of the scanner, and the average value of the shear modulus over the region of interest was calculated (Figure 1). At each time point, the values of three measurements in each trial were averaged and used for further analyses.




Figure 1. Typical examples of ultrasound shear wave elastography measurements of the biceps femoris long head (BFlh) in concentric (CON) and eccentric (ECC) tasks. Measurements were performed before (PRE) and 30, 60, 90, and 120 s after each task (POST30, POST60, POST90, POST120, respectively). The colored region represents the shear modulus map with the scale below the images. Scale bar = 1 cm.

Before the above experiments commenced, to evaluate the inter-trial (intra-day, intra-rater) repeatability, passive BFlh shear modulus measurements were performed for both legs of 2 participants (i.e., total 4 legs) in the sitting position as mentioned above (i.e., the hip flexed at 70° and knee of the tested leg fully extended). Measurements were performed 4 times for each leg, with a 2-min rest between measurements. During the rest period, participants were allowed to flex their legs.

Statistical analysis

The intraclass correlation coefficient (ICC) and coefficient of variation (CV) were calculated to evaluate the repeatability. For the main experimental data, the normality of the data was confirmed using the Shapiro–Wilk test. A two-way analysis of variance (ANOVA) [TASK (CON, ECC) × TIME (PRE and POST30, POST60, POST90, and POST120)] with repeated measures was performed. When a significant interaction was identified, additional one-way ANOVA with post-hoc tests (Dunnnett test and paired t-test) were performed. The significance level for all comparisons was set at $p = 0.05$. All statistical analyses were performed using statistical software (Prism, version 9.3.1; GraphPad Software). Data are expressed as mean \pm SD.

Results

For the inter-trial repeatability, the CV was $\leq 2.7\%$, with an ICC of 0.94, indicating excellent inter-trial repeatability.




Figure 2. Passive biceps femoris long head (BFlh) shear modulus before (PRE) and 30, 60, 90, and 120 s after (POST30, POST60, POST90, POST120, respectively) concentric (CON) and eccentric (ECC) exercises. *Significant reduction compared with PRE in ECC task ($p < 0.05$).

Figure 2 shows the passive BFlh shear modulus in CON and ECC tasks. Two-way ANOVA revealed a significant TASK × TIME interaction ($p = 0.012$, partial $\eta^2 = 0.372$). Post-hoc tests showed that passive BFlh shear modulus in ECC task was significantly reduced at POST30 ($p = 0.023$, Cohen's $d = 0.95$), POST60 ($p = 0.004$, Cohen's $d = 1.30$), POST90 ($p = 0.002$, Cohen's $d = 1.39$) and

POST120 ($p = 0.001$, Cohen's $d = 1.44$) compared with PRE while there was no significant main effect of TIME on passive BFlh shear modulus in CON task ($p = 0.473$, partial $\eta^2 = 0.062$).

Discussion

The finding of the present study was that the passive BFlh shear modulus was reduced after low-volume, slow-velocity ECC while there was no significant change in the CON task. The results support our hypothesis that eccentric knee flexion could acutely reduce passive BFlh stiffness. To the best of our knowledge, this is the first study to demonstrate contraction mode-dependent acute changes in passive muscle stiffness by using ultrasound SWE.

Passive muscle stiffness has been reported to reduce after an acute bout of passive lengthening (i.e., passive stretching exercise) (Hirata et al., 2017; Hirata et al., 2016; Miyamoto et al., 2017). Although several possible mechanisms have been proposed for the passive lengthening-induced reduction in passive muscle stiffness (Kay et al., 2015; Nakamura et al., 2011; Nakamura et al., 2015), the most likely and plausible explanation is intramuscular connective tissue lengthening (Gajdosik, 2001; Morse et al., 2008). Intramuscular connective tissues such as perimysium and endomysium have wavy fibrils, which are commonly known as crimp (Diamant et al., 1972). Previous studies have shown that the crimps within the connective tissues disappeared after lengthening the tissues (Franchi et al., 2007b). Furthermore, it is suggested that such crimps decrease in number, appear more flattened, and do not reappear after the tissue is physiologically elongated to an extent in vivo (Franchi et al., 2007a; Franchi et al., 2007b). It is reasonable to presume that these phenomena would occur during passive lengthening and during active lengthening (i.e., eccentric exercise).

Another possible mechanism for reduced passive BFlh stiffness is the increased muscle temperature (Sapin de Brosses et al., 2010). In the present study, five repetitions of 4.5 s (calculated based on 90° range of motion and angular velocity of 20°/s) isokinetic knee flexion were performed with maximal efforts in both CON and ECC tasks. Thus, it is most unlikely that muscle temperature was elevated only in the ECC task, but not in the CON task, although the muscle temperature in BFlh has not been assessed.

The effect of static stretching duration on passive muscle stiffness has been examined using SWE in previous literature (Reiner et al., 2022; Sato et al., 2020). Reiner et al. (Reiner et al., 2022) showed that passive semitendinosus stiffness was significantly reduced after a 120-s static stretching. Also, Sato et al. (Sato et al., 2020) showed no significant change in passive medial gastrocnemius stiffness immediately after a 20-s static stretching. These findings together indicate that passive lengthening with a duration as short as 20 s could not reduce passive muscle stiffness. In contrast, the present study demonstrated that passive BFlh stiffness was significantly reduced after 5 repetitions of 4.5 s (i.e., 22.5 s in total) active lengthening. However, we should note that the procedure used in the present study included not only five consecutive repetitions

of CON or ECC knee flexion, but also a warm-up exercise between the PRE and POST measurements, although a 3-min rest period was allocated after the warm-up exercise. Even when considering the effect of the specific warm-up exercise, the duration of eccentric contractions is less than 60 s in total. Although this concern needs to be kept in mind, it is likely that active lengthening (i.e., eccentric exercise) is preferred over passive lengthening (i.e., passive stretching exercise) to reduce passive muscle stiffness more efficiently in a shorter time.

Previous studies have reported that passive muscle stiffness increased after eccentric exercises aiming to induce muscle damage and/or delayed onset muscle soreness (Leung et al., 2017; Matsuo et al., 2015; Xu et al., 2019). This is in contrast with our finding that passive BFlh stiffness was reduced in ECC task. The contradiction between the present and previous studies is most likely due to the difference in the number of repetitions and set of eccentric exercises. In the previous studies, the volumes of eccentric exercises were relatively high (15 repetitions of heel drops \times 10 sets (Leung et al., 2017), 10 maximal eccentric contractions \times 6 sets (Matsuo et al., 2015), and 75 maximal eccentric contractions \times 1 set (Xu et al., 2019)). In contrast, the eccentric exercise protocol in the present study was determined based on injury prevention warm-up programs including the Nordic hamstring exercise (e.g., FIFA 11+, Part 2-9, HAMSTRINGS for 'beginner': 1 set (3-5 repetitions) (Soligard et al., 2008)), and the volume (5 maximal isokinetic eccentric contractions \times 1 set) was considerably lower than that of the previous studies. Although the threshold for eccentric exercise volume that stiffens the exercising muscle is unknown at this time, a well-defined threshold volume should be crucial to injury prevention programs. Further studies are required to clarify the relationship between eccentric exercise volume and subsequent change in muscle stiffness.

The effectiveness of eccentric hamstring exercise training (e.g., hamstring training intervention involving the Nordic hamstring exercise) for the prevention of HSI has been demonstrated (van Dyk et al., 2019). Although an increase in BFlh fascicle length induced by eccentric hamstring training has been suggested as a possible mechanism for the prevention of HSI, available information is limited on the favorable adaptive responses in the BFlh to eccentric hamstring training, and it remains to be elucidated how eccentric hamstring training influences the BFlh. The current findings that low-volume, slow-velocity eccentric knee flexion could acutely reduce passive BFlh stiffness, which is suggested as a potential risk factor for the HSI although this has not been epidemiologically demonstrated (Kumagai et al., 2019; Miyamoto-Mikami et al., 2021), will be evidence for the background of argument regarding the protective effect of eccentric hamstring training on the HSI. Longitudinal studies with eccentric hamstring training intervention are warranted to examine whether low-volume, slow-velocity eccentric hamstring training can simultaneously reduce passive BFlh stiffness and increase BFlh fascicle length and which exercises are more effective in concurrently reducing passive BFlh stiffness and lengthening BFlh fascicles.

The present study has some limitations. First, the

passive BFlh shear modulus was measured only at 50% of thigh length. Kellis et al. (Kellis et al., 2010) reported that the biceps femoris architecture (e.g., fascicle length) is not uniform from proximal to distal positions. This previous finding suggests a possibility that mechanical stress imposed during passive lengthening (passive stretching exercise) is nonuniform within a muscle (Miyamoto et al., 2017), which may lead to inhomogeneity in muscle stiffness changes. Besides, Hegyi et al. (Hegyi et al., 2019) showed the proximal-distal differences in electromyographic activity of BFlh during common hamstrings exercises. Although it remains unknown whether such proximal-distal differences in electromyographic activity can lead to the regional difference in muscle stiffness changes at this time, the change in passive muscle stiffness after active lengthening (eccentric exercise) may be nonuniform along the BFlh. Further studies are required to reveal whether the current findings hold true for other regions. Second, we failed to observe the return to baseline, although passive BFlh stiffness was measured every 30 s until 2 min after the end of eccentric exercise. Prior to the study, because there are no studies on the time course of acute changes in passive muscle stiffness after low-volume eccentric contraction, we expected that the acute effect of eccentric contractions of duration as short as 22.5 s (5 repetitions of 4.5 s) would disappear within 2 min, based on the previous findings that the effect of static stretching of duration on passive muscle stiffness (Kay and Blazevich, 2009; Konrad et al., 2019; Konrad and Tilp, 2020). Additionally, in the present study, the acute effect of eccentric contractions was only examined at the angular velocity of 20°/s. Therefore, future studies with larger sample size are warranted to examine how long the eccentric exercise-induced acute reduction in passive muscle stiffness can last, with special emphasis on eccentric exercise volume (repetition, set, intensity, and/or duration) and angular velocity.

Conclusion

The present study demonstrated that passive BFlh stiffness was acutely reduced after five repetitions of maximal isokinetic eccentric but not concentric knee flexion, suggesting that passive BFlh stiffness would reduce specifically after low-volume, slow-velocity eccentric knee flexion exercise. The present findings may provide practitioners with a basis to develop more effective exercise programs for preventing HSI.

Acknowledgements

The authors would like to thank all the students for their participation in the present study and the school where the study took place. The experiments comply with the current laws of the country in which they were performed. This work was supported in part by the Institute of Health and Sports Science & Medicine, Juntendo University. The authors have no conflicts of interest to declare. The datasets generated during and/or analyzed during the current study are not publicly available, but are available from the corresponding author who was an organizer of the study.

References

- Ahmad, C.S., Dick, R.W., Snell, E., Kenney, N.D., Curriero, F.C., Pollack, K., Albright, J.P. and Mandelbaum, B.R. (2014) Major and minor league baseball hamstring injuries: Epidemiologic

- findings from the major league baseball injury surveillance system. *American Journal of Sports Medicine* **42**, 1464-1470. <https://doi.org/10.1177/0363546514529083>
- Alt, T., Nodler, Y.T., Severin, J., Knicker, A.J. and Strüder, H.K. (2018) Velocity-specific and time-dependent adaptations following a standardized Nordic hamstring exercise training. *Scandinavian Journal of Medicine & Science in Sports* **28**, 65-76. <https://doi.org/10.1111/sms.12868>
- Askling, C.M., Tengvar, M., Saartok, T. and Thorstensson, A. (2007) Acute first-time hamstring strains during high-speed running: A longitudinal study including clinical and magnetic resonance imaging findings. *American Journal of Sports Medicine* **35**, 197-206. <https://doi.org/10.1177/0363546506294679>
- Brooks, J.H., Fuller, C.W., Kemp, S.P. and Reddin, D.B. (2005) Epidemiology of injuries in English professional rugby union: part 1 match injuries. *British Journal of Sports Medicine* **39**, 757-766. <https://doi.org/10.1136/bjsm.2005.018135>
- Diamant, J., Keller, A., Baer, E., Litt, M. and Arridge, R.G. (1972) Collagen; ultrastructure and its relation to mechanical properties as a function of ageing. *Proceedings of the Royal Society of London. Series B. Biological Sciences* **180**, 293-315. <https://doi.org/10.1098/rspb.1972.0019>
- Ekstrand, J., Hagglund, M. and Walden, M. (2011) Epidemiology of muscle injuries in professional football (soccer). *American Journal of Sports Medicine* **39**, 1226-1232. <https://doi.org/10.1177/0363546510395879>
- El-Ashker, S., Allardyce, J.M. and Carson, B.P. (2019) Sex-related differences in joint-angle-specific hamstring-to-quadriceps function following fatigue. *European Journal of Sport Science* **19**, 1053-1061. <https://doi.org/10.1080/17461391.2019.1574904>
- Franchi, M., Fini, M., Quaranta, M., De Pasquale, V., Raspanti, M., Giavaresi, G., Ottani, V. and Ruggeri, A. (2007a) Crimp morphology in relaxed and stretched rat Achilles tendon. *Journal of Anatomy* **210**, 1-7. <https://doi.org/10.1111/j.1469-7580.2006.00666.x>
- Franchi, M., Trire, A., Quaranta, M., Orsini, E. and Ottani, V. (2007b) Collagen structure of tendon relates to function. *The Scientific World Journal* **7**, 404-420. <https://doi.org/10.1100/tsw.2007.92>
- Gajdosik, R.L. (2001) Passive extensibility of skeletal muscle: Review of the literature with clinical implications. *Clinical Biomechanics* **16**, 87-101. [https://doi.org/10.1016/s0268-0033\(00\)00061-9](https://doi.org/10.1016/s0268-0033(00)00061-9)
- Gerard, R., Gojon, L., Declève, P. and Van Cant, J. (2020) The effects of eccentric training on biceps femoris architecture and strength: A systematic review with meta-analysis. *Journal of Athletic Training* **55**, 501-514. <https://doi.org/10.4085/1062-6050-194-19>
- Green, B., Bourne, M.N., van Dyk, N. and Pizzari, T. (2020) Recalibrating the risk of hamstring strain injury (HSI): A 2020 systematic review and meta-analysis of risk factors for index and recurrent hamstring strain injury in sport. *British Journal of Sports Medicine* **54**, 1081-1088. <https://doi.org/10.1136/bjsports-2019-100983>
- Hegyí, A., Csala, D., Peter, A., Finni, T. and Cronin, N.J. (2019) High-density electromyography activity in various hamstring exercises. *Scandinavian Journal of Medicine & Science in Sports* **29**, 34-43. <https://doi.org/10.1111/sms.13303>
- Hirata, K., Kanehisa, H. and Miyamoto, N. (2017) Acute effect of static stretching on passive stiffness of the human gastrocnemius fascicle measured by ultrasound shear wave elastography. *European Journal of Applied Physiology* **117**, 493-499. <https://doi.org/10.1007/s00421-017-3550-z>
- Hirata, K., Miyamoto-Mikami, E., Kanehisa, H. and Miyamoto, N. (2016) Muscle-specific acute changes in passive stiffness of human triceps surae after stretching. *European Journal of Applied Physiology* **116**, 911-918. <https://doi.org/10.1007/s00421-016-3349-3>
- Kay, A.D. and Blazevich, A.J. (2009) Moderate-duration static stretch reduces active and passive plantar flexor moment but not Achilles tendon stiffness or active muscle length. *Journal of Applied Physiology* **106**, 1249-1256. <https://doi.org/10.1152/jappphysiol.91476.2008>
- Kay, A.D., Husbands-Beasley, J. and Blazevich, A.J. (2015) Effects of contract-relax, static stretching, and isometric contractions on muscle-tendon mechanics. *Medicine and Science in Sports and Exercise* **47**, 2181-2190. <https://doi.org/10.1249/MSS.0000000000000632>
- Kellis, E., Galanis, N., Natsis, K. and Kapetanios, G. (2010) Muscle architecture variations along the human semitendinosus and biceps femoris (long head) length. *Journal of Electromyography and Kinesiology* **20**, 1237-1243. <https://doi.org/10.1016/j.jelekin.2010.07.012>
- Konrad, A., Reiner, M.M., Thaller, S. and Tilp, M. (2019) The time course of muscle-tendon properties and function responses of a five-minute static stretching exercise. *European Journal of Sport Science* **19**, 1195-1203. <https://doi.org/10.1080/17461391.2019.1580319>
- Konrad, A. and Tilp, M. (2020) The time course of muscle-tendon unit function and structure following three minutes of static stretching. *Journal of Sports Science and Medicine* **19**, 52-58.
- Kumagai, H., Miyamoto-Mikami, E., Hirata, K., Kikuchi, N., Kamiya, N., Hoshikawa, S., Zempo, H., Naito, H., Miyamoto, N. and Fuku, N. (2019) ESR1 rs2234693 polymorphism is associated with muscle injury and muscle stiffness. *Medicine and Science in Sports and Exercise* **51**, 19-26. <https://doi.org/10.1249/MSS.0000000000001750>
- Leung, W.K.C., Chu, K.L. and Lai, C. (2017) Sonographic evaluation of the immediate effects of eccentric heel drop exercise on Achilles tendon and gastrocnemius muscle stiffness using shear wave elastography. *PeerJ* **5**, e3592. <https://doi.org/10.7717/peerj.3592>
- Matsuo, S., Suzuki, S., Iwata, M., Banno, Y., Asai, Y., Tsuchida, W. and Inoue, T. (2013) Acute effects of different stretching durations on passive torque, mobility, and isometric muscle force. *The Journal of Strength & Conditioning Research* **27**, 3367-3376. <https://doi.org/10.1519/JSC.0b013e318290c26f>
- Matsuo, S., Suzuki, S., Iwata, M., Hatano, G. and Nosaka, K. (2015) Changes in force and stiffness after static stretching of eccentrically-damaged hamstrings. *European Journal of Applied Physiology* **115**, 981-991. <https://doi.org/10.1007/s00421-014-3079-3>
- Miyamoto-Mikami, E., Kumagai, H., Tanisawa, K., Taga, Y., Hirata, K., Kikuchi, N., Kamiya, N., Kawakami, R., Midorikawa, T., Kawamura, T., Kakigi, R., Natsume, T., Zempo, H., Suzuki, K., Kohmura, Y., Mizuno, K., Torii, S., Sakamoto, S., Oka, K., Higuchi, M., Naito, H., Miyamoto, N. and Fuku, N. (2021) Female athletes genetically susceptible to fatigue fracture are resistant to muscle injury: Potential role of COL1A1 variant. *Medicine and Science in Sports and Exercise* **53**, 1855-1864. <https://doi.org/10.1249/MSS.00000000000002658>
- Miyamoto, N., Hirata, K. and Kanehisa, H. (2017) Effects of hamstring stretching on passive muscle stiffness vary between hip flexion and knee extension maneuvers. *Scandinavian Journal of Medicine & Science in Sports* **27**, 99-106. <https://doi.org/10.1111/sms.12620>
- Miyamoto, N., Hirata, K., Kimura, N. and Miyamoto-Mikami, E. (2018) Contributions of hamstring stiffness to straight-leg-raise and sit-and-reach test scores. *International Journal of Sports Medicine* **39**, 1108-1114. <https://doi.org/10.1055/s-0043-117411>
- Mjolsnes, R., Arnason, A., Osthagen, T., Raastad, T. and Bahr, R. (2004) A 10-week randomized trial comparing eccentric vs. concentric hamstring strength training in well-trained soccer players. *Scandinavian Journal of Medicine & Science in Sports* **14**, 311-317. <https://doi.org/10.1046/j.1600-0838.2003.367.x>
- Morse, C.I., Degens, H., Seynnes, O.R., Manganaris, C.N. and Jones, D.A. (2008) The acute effect of stretching on the passive stiffness of the human gastrocnemius muscle tendon unit. *The Journal of Physiology* **586**, 97-106. <https://doi.org/10.1113/jphysiol.2007.140434>
- Nakamura, M., Ikezoe, T., Takeno, Y. and Ichihashi, N. (2011) Acute and prolonged effect of static stretching on the passive stiffness of the human gastrocnemius muscle tendon unit in vivo. *Journal of Orthopaedic Research* **29**, 1759-1763. <https://doi.org/10.1002/jor.21445>
- Nakamura, M., Ikezoe, T., Takeno, Y. and Ichihashi, N. (2013) Time course of changes in passive properties of the gastrocnemius muscle-tendon unit during 5 min of static stretching. *Manual Therapy* **18**, 211-215. <https://doi.org/10.1016/j.math.2012.09.010>
- Nakamura, M., Ikezoe, T., Tokugawa, T. and Ichihashi, N. (2015) Acute effects of stretching on passive properties of human gastrocnemius muscle-tendon unit: Analysis of differences between hold-relax and static stretching. *Journal of Sport Rehabilitation* **24**, 286-292. <https://doi.org/10.1123/jsr.2014-0164>

- O'Sullivan, K., McAuliffe, S. and Deburca, N. (2012) The effects of eccentric training on lower limb flexibility: A systematic review. *British Journal of Sports Medicine* **46**, 838-845. <https://doi.org/10.1136/bjsports-2011-090835>
- Opar, D.A., Drezner, J., Shield, A., Williams, M., Webner, D., Sennett, B., Kapur, R., Cohen, M., Ulager, J., Cafengiu, A. and Cronholm, P.F. (2014) Acute hamstring strain injury in track-and-field athletes: A 3-year observational study at the Penn Relay Carnival. *Scandinavian Journal of Medicine & Science in Sports* **24**, e254-e259. <https://doi.org/10.1111/sms.12159>
- Reiner, M.M., Tilp, M., Guilhem, G., Morales-Artacho, A. and Konrad, A. (2022) Comparison of A Single Vibration Foam Rolling and Static Stretching Exercise on the Muscle Function and Mechanical Properties of the Hamstring Muscles. *Journal of Sports Science and Medicine* **21**, 287-297. <https://doi.org/10.52082/jssm.2022.287>
- Sapin-de Brosse, E., Gennisson, J.L., Pernot, M., Fink, M. and Tanter, M. (2010) Temperature dependence of the shear modulus of soft tissues assessed by ultrasound. *Physics in Medicine and Biology* **55**, 1701-1718. <https://doi.org/10.1088/0031-9155/55/6/011>
- Sato, S., Kiyono, R., Takahashi, N., Yoshida, T., Takeuchi, K. and Nakamura, M. (2020) The acute and prolonged effects of 20-s static stretching on muscle strength and shear elastic modulus. *PLoS One* **15**, e0228583. <https://doi.org/10.1371/journal.pone.0228583>
- Severo-Silveira, L., Dornelles, M.P., Lima, E.S.F.X., Marchiori, C.L., Medeiros, T.M., Pappas, E. and Baroni, B.M. (2021) Progressive workload periodization maximizes effects of Nordic hamstring exercise on muscle injury risk factors. *The Journal of Strength & Conditioning Research* **35**, 1006-1013. <https://doi.org/10.1519/JSC.0000000000002849>
- Soligard, T., Myklebust, G., Steffen, K., Holme, I., Silvers, H., Bizzini, M., Junge, A., Dvorak, J., Bahr, R. and Andersen, T.E. (2008) Comprehensive warm-up programme to prevent injuries in young female footballers: Cluster randomised controlled trial. *British Medical Journal* **337**, a2469. <https://doi.org/10.1136/bmj.a2469>
- Timmins, R.G., Bourne, M.N., Shield, A.J., Williams, M.D., Lorenzen, C. and Opar, D.A. (2016) Short biceps femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite football (soccer): a prospective cohort study. *British Journal of Sports Medicine* **50**, 1524-1535. <https://doi.org/10.1136/bjsports-2015-095362>
- van Dyk, N., Behan, F.P. and Whiteley, R. (2019) Including the Nordic hamstring exercise in injury prevention programmes halves the rate of hamstring injuries: A systematic review and meta-analysis of 8459 athletes. *British Journal of Sports Medicine* **53**, 1362-1370. <https://doi.org/10.1136/bjsports-2018-100045>
- Watsford, M.L., Murphy, A.J., McLachlan, K.A., Bryant, A.L., Cameron, M.L., Crossley, K.M. and Makkdissi, M. (2010) A prospective study of the relationship between lower body stiffness and hamstring injury in professional Australian rules footballers. *American Journal of Sports Medicine* **38**, 2058-2064. <https://doi.org/10.1177/0363546510370197>
- Xu, J., Fu, S.N., Zhou, D., Huang, C. and Hug, F. (2019) Relationship between pre-exercise muscle stiffness and muscle damage induced by eccentric exercise. *European Journal of Sport Science* **19**, 508-516. <https://doi.org/10.1080/17461391.2018.1535625>

Key points

- Passive muscle stiffness of the biceps femoris long head (BFlh) was directly assessed using ultrasound shear wave elastography.
- Passive BFlh stiffness was reduced after five consecutive repetitions of eccentric knee flexion, but not concentric contractions.
- The present findings may provide practitioners with a basis to develop more effective exercise programs for preventing HSI.

AUTHOR BIOGRAPHY



Lei ZHI

Employment

Graduate School of Health and Sports Science, Juntendo University, Inzai, Japan.

Degree

MDResearch interests

Exercise physiology, Sports Medicine.

E-mail: jurgenzi@gmail.com



Naokazu MIYAMOTO

Employment

Graduate School of Health and Sports Science, Juntendo University, Inzai, Japan.

Degree

PhD

Research interest

Neuromuscular physiology, Biomechanics.

E-mail: n-miyamoto@juntendo.ac.jp



Hisashi NAITO

Employment

Graduate School of Health and Sports Science, Juntendo University, Inzai, Japan.

Degree

PhD

Research interest

Exercise physiology.

E-mail: hnaitou@juntendo.ac.jp

✉ Naokazu Miyamoto

Faculty of Health and Sports Science, Juntendo University 1-1 Hiraka-gakuendai, Inzai, Chiba 270-1695, Japan